THE CANADIAN ENTOMOLOGIST

Index to Volume 133 (2001)

THE ENTOMOLOGICAL SOCIETY OF CANADA, OTTAWA

Acar EB, Medina JC., Lee ML, Booth GM. Olfactory behavior of convergent lady beetles (Coleoptera: Coccinellidae) to alarm pheromone of green peach aphid (Hemiptera: Aphididae), 389
Alarie Y, Watts CHS, Nilsson AN. Larval morphology of the tribe Matini (Coleoptera: Dytiscidae, Colymbetinae): descriptions of *Batrachomatus daemeli*, *Matus bicarinatus*, and *Allomatus nannup* and phylogenetic relationships, 165
Amezaga I, Garbisu C. Effect of maturation feeding period on survival of *Tomicus piniperda* (Coleoptera: Scolytidae), 131
Andersen A, McNeil JN. Size influences male mating success in the alfalfa blotch leafminer (Diptera: Agromyzidae), 717
Anderton LK, Jenkins MJ. Cone entomofauna of whitebark pine and alpine larch (Pinaceae): potential impact of *Leptoglossus occidentalis* (Hemiptera: Coreidae) and a new record of *Strobilomyia macalpinei* (Diptera: Anthomyiidae), 399
Archangelsky M, Branham MA. Description of last instar and pupa of *Pyropyga nigricans* (Coleoptera: Lampyridae, Photinini) and comparison with larvae of other Photinini genera, 155
Baldwin D. See Braun *et al.*, 565
Bates SL. See Strong *et al.*, 857
Bauce É. See Trudel *et al.*, 333
Behan-Pelletier VM, Eamer B, Clayton M. Mycobatidae (Acari: Oribatida) of Pacific Northwest canopy habitats, 755
Bentz BJ, Logan JA, Vandygriff JC. Latitudinal variation in *Dendroctonus ponderosae* (Coleoptera: Scolytidae) development time and adult size, 375
See Hansen *et al.*, 805, 827
Berleur G, Gingras J, Tourneur J-C. Influence of diet on development and oviposition of *Forficula auricularia* (Dermaptera: Forficulidae), 705
Berthiaume R, Hébert C, Cloutier C. *Podabrus rugosulus* (Coleoptera: Cantharidae), an opportunist predator of *Mindarus abietinus* (Hemiptera: Aphididae) in Christmas tree plantations, 151
Biondi M. New flea beetle genus and species (Coleoptera: Chrysomelidae, Alticinae) from Central Africa, 643
Bittner LA. See Pree *et al.*, 93
Booth GM. See Acar *et al.*, 389
Boivin G. See Roitberg *et al.*, 429
Borden JH. See Broberg *et al.*, 119
Boucher S, Wheeler TA. Diversity of Agromyzidae (Diptera) in disjunct grasslands of the southern Yukon Territory, 593
Brailovsky H. New genera and species of the tribe Acanthocorini (Hemiptera: Coreidae, Coreinae), 623
Branham MA. See Archangelsky and Branham, 155
Braun L, Erlandson M, Baldwin D, Soroka J, Mason P, Foottit R, Hegedus D. Seasonal occurrence, species composition, and parasitism of *Lygus* spp. in alfalfa, canola, and mustard, 565
Broberg CL, Borden JH, Humble LM. Host range, attack dynamics, and impact of *Cryptorhynchus lapathi* (Coleoptera: Curculionidae) on *Salix* (Salicaceae) spp., 119
Brodeur J, Leclerc L-A, Fournier M, Roy M. Cabbage seedpod weevil (Coleoptera: Curculionidae): new pest of canola in northeastern North America, 709
Buck M. Protogyny, protandry, and bimodal emergence patterns in necrophagous Diptera, 521
Carlson JE. See Lewis *et al.*, 229
Casagrande RA. See Gold *et al.*, 671
Castrezana S, Markow TA. Arthropod diversity in necrotic tissue of three species of columnar cacti (Cactaceae), 301
Castro D del C, Cicchino A. New species of the genus *Phtheiropoios* (Phthiraptera: Gyropidae), parasitic on *Ctenomys* (Rodentia: Octodontidae), 87
Chandler LD. See Mulock and Chandler, 105
Chau A, Mackauer M. Host–instar selection in the aphid parasitoid *Monoctonus paulensis* (Hymenoptera: Braconidae, Aphidiinae): assessing costs and benefits, 549
Chiew PK. See Rhainds *et al.*, 409
Christianson BA. See Knight *et al.*, 413
Cicchino AC. See Castro D del and Cicchino, 87
See Roig-Juñent and Cicchino, 651
Clarke J. See Lamb *et al.*, 579
Clarke P. See Lamb *et al.*, 579
Clayton M. See Behan-Pelletier *et al.*, 755
Cloutier C. See Berthiaume *et al.*, 151
Cooper BE. See Vakenti *et al.*, 139
Cossentine JE. See Vakenti *et al.*, 139
Dotseth EJ, Johnson NF. Revision of the Neotropical genus *Acanthoscelio* (Hymenoptera: Scelionidae), 487
Eamer B. See Behan-Pelletier *et al.*, 755

El-Kassaby YA. See Lewis *et al.*, 229
El-Sayed AM, Fraser HM, Trimble RM. Chemical identification and behavioral activity of (*Z*)-11-tetradecenal in an eastern North American population of the obliquebanded leafroller, 365
Modification of the sex-pheromone communication system associated with organophosphorus-insecticide resistance in the obliquebanded leafroller (Lepidoptera: Tortricidae), 867
Erlandson M. See Braun *et al.*, 565
See Ulmer *et al.*, 509
Eveleigh ES. See Piene *et al.*, 675
Fields P. See Hou *et al.*, 325
Fielding DJ, Brusven MA, Shafii B, Price WJ. Spatial heterogeneity of low-density populations of *Melanoplus sanguinipes* (Orthoptera: Acrididae) associated with grazing and vegetation treatments, 843
Floate K. See Gibson and Floate, 49
Foottit R. See Braun *et al.*, 565
Fournier M. See Brodeur *et al.*, 709
Fraser HM. See El-Sayed *et al.*, 365, 867
Fraser HW, Trimble RM. Effect of delayed mating on reproductive biology of the Oriental fruit moth (Lepidoptera: Tortricidae), 219
Freitag R, Hartwick A, Singh A. Flagellar microstructures of male tiger beetles (Coleoptera: Cicindelidae): implications for systematics and functional morphology, 633
Galloway T. See Hou *et al.*, 325
Garbisu C. See Amezaga and Garbisu, 131
Génier F. See Smith and Génier, 777
Germain C. See Klimaszewski *et al.*, 1
Gibson GAP, Floate K. Species of *Trichomalopsis* (Hymenoptera: Pteromalidae) associated with filth flies (Diptera: Muscidae) in North America, 49
Gillott C. See Ulmer *et al.*, 509
Gingras J. See Berleur *et al.*, 705
Gingras J, Tourneur J-C. Timing of adult mortality, oviposition, and hatching during the underground phase of *Forficula auricularia* (Dermaptera: Forficulidae), 269
Glaubitz J. See Lewis *et al.*, 229
Glenn DM. See Knight *et al.*, 413
Glubish SS. See Reid and Glubish, 697
Gold MS, Casagrande RA, Tewksbury LA, Livingston SB, Kenis M. European parasitoids of *Lilioceris lilii* (Coleoptera: Chrysomelidae), 671
Gries G. See Takács and Gries, 725
See Rhainds *et al.*, 409
Hallett RH, Heal JD. First Nearctic record of the swede midge (Diptera: Cecidomyiidae), a pest of cruciferous crops from Europe, 713
Hansen EM, Bentz BJ, Turner DL. Physiological basis for flexible voltinism in the spruce beetle (Coleoptera: Scolytidae), 805
Temperature-based model for predicting univoltine brood proportionas in spruce bettle (Coleoptera: Scolytidae), 827
Hartwick A. See Freitag *et al.*, 633
Heal JD. See Hallett and Heal, 713
Hébert C. See Berthiaume *et al.*, 151
See Klimaszewski *et al.*, 1
Hegedus D. See Braun *et al.*, 565
Helson BV, Lyons DB, Wanner KW, Scarr TA. Control of conifer defoliators with neem-based systemic bioinsecticides using a novel injection device, 729
Hinojosa T. See Horton *et al.*, 109
Hollmann J. See Sahota *et al.*, 315
Horton DR, Lewis TM, Hinojosa T. Copulation duration and probability of insemination in *Anthocoris whitei* (Hemiptera: Anthocoridae) as a function of male body size, 109
Hou X, Fields P, Galloway T. Cold hardiness of black carpet beetle larvae, 325
Humble LM. See Broberg *et al.*, 119
See Klimaszewski *et al.*, 1
Huang D-Y. See Lin and Huang, 747
Hurley RL. See Runyon *et al.*, 215
Ibaraki A. See Sahota *et al.*, 315
Inouye DW. See Larson *et al.*, 439
Jenkins MJ. See Anderton and Jenkins, 399
Jensen LBM. See Vakenti *et al.*, 139
Johnson NF. See Dotseth and Johnson, 487

Kenis M. See Gold *et al.*, 671
Kevan PG. See Larson *et al.*, 439
See Morandin *et al.*, 883
Khosla S. See Morandin *et al.*, 883
Klimaszewski J, Pelletier G, Germain C, Hébert C, Humble LM, Winchester NN. Diversity of *Placusa* (Coleoptera: Staphylinidae, Aleocharinae) in Canada, with descriptions of two new species, 1
Knight AL, Christianson BA, Unruh TR, Puterka G, Glenn DM. Impacts of seasonal kaolin particle films on apple pest management, 413
Kuenen LPS, Silk PJ. Sex-biased trap capture and odor-stimulated upwind flight in the field by *Rhagoletis mendax* (Diptera: Tephritidae), 293
Kuhlmann U. See Vincent *et al.*, 147
Lamb RJ. See Smith and Lamb, 533
See Wise *et al.*, 255
Lamb RJ, Smith MAH, Wise IL, Clarke P, Clarke J. Oviposition deterrence to *Sitodiplosis mosellana* (Diptera: Cecidomyiidae): a source of resistance for durum wheat (Gramineae), 579
Lan CC. See Rhainds *et al.*, 409
Larson BMH, Kevan PG, Inouye DW. Flies and flowers: taxonomic diversity of anthophiles and pollinators, 439
Lavallée R. See Trudel *et al.*, 333
Laverty TM. See Morandin *et al.*, 883
Leal I. See Sahota *et al.*, 315
Leclerc L-A. See Brodeur *et al.*, 709
Lee ML. See Acar *et al.*, 389
Lester PJ. See Marshall *et al.*, 279
Lewis KG, Ritland K, El-Kassaby YA, McLean JA, Glaubitz J, Carlson JE. Randomly amplified polymorphic DNA reveals fine-scale genetic structure in *Pissodes strobi* (Coleoptera: Curculionidae), 229
Lewis TM. See Horton *et al.*, 109
Li SY, Otvos IS. Differences in activity enhancement of a baculovirus by an optical brightener between *Choristoneura fumiferana* and *Choristoneura occidentalis* (Lepidoptera: Tortricidae), 329
Lin Q-B, Huang D-Y. Description of *Caenoephemera shangyuanensis*, **gen.nov., sp.nov.** (Ephemeroptera), from the Yixian Formation, 747
Livingston SB. See Gold *et al.*, 671
Logan JA. See Bentz *et al.*, 375
Lyons DB. See Helson *et al.*, 729
Mackauer M. See Chau and Mackauer, 549
Manville JF. See Sahota *et al.*, 315
Markow TA. See Castrezana and Markow, 301
Marshall DB, Thistlewood HMA, Lester PJ. Release, establishment, and movement of the predator *Typhlodromus pyri* (Acari: Phytoseiidae) on apple, 279
See Pogoda *et al.*, 819
Mason P. See Braun *et al.*, 565
McCullough DG. See Siegert and McCullough, 343
McLean JA. See Lewis *et al.*, 229
McNeil JN. See Andersen and McNeil, 717
Medina JC. See Acar *et al.*, 389
Miller DR. Frontalin interrupts attraction of *Ips pini* (Coleoptera: Scolytidae) to ipsdienol, 407
Morandin LA, Laverty TM, Kevan PG, Khosla S, Shipp L. Bumble bee (Hymenoptera: Apidae) activity and loss in commercial tomato greenhouses, 883
Morrill WL. See Runyon *et al.*, 215
Mulock BS, Chandler LD. *Beauveria bassiana* transmission in western corn rootworm (Coleoptera: Chrysomelidae), 105
Nilsson AN. See Alarie *et al.*, 165
Nosil P. Sexual size dimorphism in a natural population of *Callicorixa vulnerata* (Hemiptera: Corixidae), 311
Nyrop JP. See Waldstein *et al.*, 687
Ostaff DP. See Piene *et al.*, 675
Otis GW. See van Engelsdorp and Otis, 793
Ötvös IS. See Szöcs *et al.*, 297
See Li and Otvos, 329
Pelletier G. See Klimaszewski *et al.*, 1
Piene H, Ostaff DP, Eveleigh ES. Growth loss and recovery following defoliation by the balsam fir sawfly in young, spaced balsam fir stands, 675
Pogoda MK, Pree DJ, Marshall DB. Effects of encapsulation on the toxicity of insecticides to the Oriental fruit moth (Lepidoptera: Tortricidae) and the predator *Typhlodromus pyri* (Acari: Phytoseiidae), 819

See Pree *et al.*, 93
Polavarapu S. See Teixeira and Polavarapu, 239
Pree DJ, Whitty KJ, Pogoda MK, Bittner LA. Occurrence of resistance to insecticides in populations of the obliquebanded leafroller from orchards, 93
See Pogoda *et al.*, 819
Prokopy RJ. See Rull and Prokopy, 355
Puterka G. See Knight *et al.*, 413
Quednau FW. World review of the genus *Tinocallis* (Hemiptera: Aphididae, Calaphidinae) with description of a new species, 197
Rancourt B. See Vincent *et al.*, 147
Raworth DA. Development, larval voracity, and greenhouse releases of *Stethorus punctillum* (Coleoptera: Coccinellidae), 721
Reid ML, Glubish SS. Tree size and growth history predict breeding densities of Douglas-fir beetles in fallen trees, 697
Reissig WH. See Waldstein *et al.*, 687
Rhainds M, Warndorff M, Chiew PK, Lan CC, Gries G. Stimuli increasing oviposition by female coffee white stem borer (Coleoptera: Cerambycidae), 409
Ritland K. See Lewis *et al.*, 229
Roig-Juñent S, Cicchino AC. *Chaltenia patagonica*, new genus and species belonging to Chalteniina, a new subtribe of Zolini (Coleoptera: Carabidae), 651
Roitberg BD, Boivin G, Vet LEM. Fitness, parasitoids, and biological control: an opinion, 429
Roy M. See Brodeur *et al.*, 709
Rull J, Prokopy RJ. Effect of apple-orchard structure on interception of *Rhagoletis pomonella* (Diptera: Tephritidae) flies by odor-baited traps, 355
Runyon JB, Hurley RL, Morrill WL, Weaver DK. Distinguishing adults of *Bracon cephi* and *Bracon lissogaster* (Hymenoptera: Braconidae), parasitoids of the wheat stem sawfly (Hymenoptera: Cephidae), 215
Sahota TS, Manville JF, Hollmann J, Leal I, Ibaraki A, White E. Resistance against *Pissodes strobi* (Coleoptera: Curculionidae) in severed leaders and in a water-soluble bark extract of *Picea sitchensis* (Pinaceae): evidence for a post-ingestive mode of action, 315
Sanders C. See Szöcs *et al.*, 297
Sarazin M. See Vincent *et al.*, 147
Siegert NW, McCullough DG. Preference of *Tomicus piniperda* (Coleoptera: Scolytidae) parent adults and shoot-feeding progeny adults for three pine species, 343
Scarr TA. See Helson *et al.*, 729
Sharkey MJ. See Vakenti *et al.*, 139
Shipp L. See Morandin *et al.*, 883
Sihvonen P. Revision of the *Scopula cajanderi* (Lepidoptera: Geometridae, Sterrhinae) species group with description of a new species, 467
Silk PJ. See Kuenen and Silk, 293
Singh A. See Freitag *et al.*, 633
Smith ABT, Génier F. Revision of the genus *Holocephalus* (Coleoptera: Scarabaeidae: Scarabaeinae: Coprini), 777
Smith MAH. See Lamb *et al.*, 579
See Wise *et al.*, 255
Smith MAH, Lamb RJ. Factors influencing oviposition by *Sitodiplosis mosellana* (Diptera: Cecidomyiidae) on wheat spikes (Gramineae), 533
Soroka J. See Braun *et al.*, 565
Stoehr MU. See Strong *et al.*, 857
Strong WB, Bates SL, Stoehr MU. Feeding by *Leptoglossus occidentalis* (Hemiptera: Coreidae) reduces seed set in lodgepole pine (Pinaceae), 857
Szöcs G, Ötvös IS, Sanders C. *Erannis tiliaria* (Lepidoptera: Geometridae) males attracted to enantiomerically identical pheromone blend of *Erannis defoliaria*, 297
Takács S, Gries G. Communication ecology of webbing clothes moth: evidence for male-produced aggregation signal(s), 725
Teixeira LAF, Polavarapu S. Occurrence of late-emerging populations of the blueberry maggot fly (Diptera: Tephritidae), 239
Tewksbury LA. See Gold *et al.*, 671
Thistlewood HMA. See Marshall *et al.*, 279
Tourneur J-C. See Berleur *et al.*, 705
See Gingras and Tourneur, 269
Trimble RM. Interaction of photoperiod and temperature in the regulation of prediapause development in the spotted tentiform leafminer (Lepidoptera: Gracillariidae), 251
See Fraser and Trimble, 219

See El-Sayed *et al.*, 365, 867
Trudel R, Lavallée R, Bauce É. Oviposition biology of *Pissodes strobi* (Coleoptera: Curculionidae) on white pine (Pinaceae) under laboratory conditions, 333
Turner DL. See Hansen *et al.*, 805, 827
Ulmer B, Gillott C, Erlandson M. Feeding preferences, growth, and development of *Mamestra configurata* (Lepidoptera: Noctuidae) on Brassicaceae, 509
Unruh TR. See Knight *et al.*, 413
van Engelsdorp D, Otis GW. Field evaluation of nine families of honey bees for resistance to tracheal mites, 793
Vakenti JM, Cossentine JE, Cooper BE, Sharkey MJ, Yoshimoto CM, Jensen LBM. Host-plant range and parasitoids of obliquebanded and three-lined leafrollers (Lepidoptera: Torticidae), 139
Vandygriff JC. See Bentz *et al.*, 375
Vet LEM. See Roitberg *et al.*, 429
Vincent C, Rancourt B, Sarazin M, Kuhlmann U. Releases and first recovery of *Lathrolestes ensator* (Hymenoptera: Ichneumonidae) in North America, a parasitoid of *Hoplocampa testudinea* (Hymenoptera: Tenthredinidae), 147
Waldstein DE, Reissig WH, Nyrop JP. Larval movement and its potential impact on the management of the obliquebanded leafroller (Lepidoptera: Tortricidae), 687
Wanner KW. See Helson *et al.*, 729
Warndorff M. See Rhainds *et al.*, 409
Watts CHS. See Alarie *et al.*, 165
Weaver DK. See Runyon *et al.*, 215
Wheeler TA. See Boucher and Wheeler, 593
White E. See Sahota *et al.*, 315
Whitty KJ. See Pree *et al.*, 93
Winchester NN. See Klimaszewski *et al.*, 1
Wise IL, Lamb RJ, Smith MAH. Domestication of wheats (Gramineae) and their susceptibility to herbivory by *Sitodiplosis mosellana* (Diptera: Cecidomyiidae), 255
Wise IL. See Lamb *et al.*, 579
Yoshimoto CM. See Vakenti *et al.*, 139

Erratum

Raworth DA. Development, larval voracity, and greenhouse releases of *Stethorus punctillum* (Coleoptera: Coccinellidae), 895